

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/319162513>

Practice of Labour Pain Management Methods and Associated Factors among Skilled Attendants Working at General Hospitals in Tigray Region, North Ethiopia: Hospital Based Cross-Secti...

Article in Health Science Journal · January 2017

DOI: 10.21767/1791-809X.1000516

CITATIONS

17

READS

1,305

7 authors, including:



Yayehyirad Yemaneh

International Medical Corps Ethiopia DolloAdo Operation

17 PUBLICATIONS 90 CITATIONS

SEE PROFILE



Addisu Alehegn Alemu

Debre Markos University

24 PUBLICATIONS 84 CITATIONS

SEE PROFILE



Salahuddinfarooq Mohammed

University of Mississippi

68 PUBLICATIONS 650 CITATIONS

SEE PROFILE



Abere Yekoye

Mekelle University

6 PUBLICATIONS 46 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Sleep, race, ethnicity and culture [View project](#)



Assesement of M aternal near miss [View project](#)

DOI: 10.21767/1791-809X.1000516

Practice of Labour Pain Management Methods and Associated Factors among Skilled Attendants Working at General Hospitals in Tigray Region, North Ethiopia: Hospital Based Cross-Sectional Study Design

Ermias Sahile^{1*}, Yayehyirad Yemaneh¹, Addisu Alehegn¹, Wendessen Nigussie², Mohhamed Salahuddin³, Abere Yekoye⁴ and Negiso Gebeyehu⁵

¹Department of Midwifery, College of Health Sciences, Mizan-Tepi University, Mizan-Teferi, Ethiopia

²Department of Nursing, College of Health Sciences, Mizan-Tepi University, Mizan-Teferi, Ethiopia

³Department of Pharmacy, College of Health Sciences, Mizan-Tepi University, Mizan-Teferi, Ethiopia

⁴Department of Midwifery, College of Health Sciences, Mekelle University, Mekelle, Ethiopia

⁵Department of Midwifery, College of Health Sciences, Wechamo university, Hosana, Ethiopia

*Corresponding author: Ermias Sahile, Lecturer, Department of Midwifery College of Health Sciences Mizan-Tepi University, Mizan-Teferi, Ethiopia, Tel: +251910110883; E-mail: ermi1380@gmail.com

Received date: 24 June 2017; Accepted date: 01 August 2017; Published date: 08 August 2017

Copyright: © 2017 Sahile E, et al. This is an open-access article distributed under the terms of the creative Commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Citation: Sahile E, Yemaneh Y, Alehegn A, Nigussie W, Salahuddin M, et al. (2017) Practice of Labour Pain Management Methods and Associated Factors among Skilled Attendants Working at General Hospitals in Tigray Region, North Ethiopia: Hospital Based Cross-Sectional Study Design. Health Sci J. Vol. 11 No. 4: 516.

Abstract

Introduction: Labour pain is the most severe pain that a woman ever has to face. Pain relief in childbirth is subject to many social and cultural modifiers, which continue to change. Delivery of the infant into the arms of a conscious and pain-free mother is the most exciting and rewarding moment in maternal care services. Developed nation give emphasis on continuous labour support, But in developing country pain is neglected especially managing labor pain. In a country like Ethiopia with low institutional delivery managing labour pain can help as an incentive to enhance institutional delivery rate.

Objective: The main purpose of this study is to assess practice of Labor pain management methods and associated factors among skilled attendants in all Tigray region general hospitals.

Method: Institutional based cross sectional study design was carried out at all general hospitals in tigray regional state from October 01-30, 2016. All skilled attendants who were working at labour ward were included in the study; Data was collected by interviewer-administered, pretested and structured questionnaire. The data were entered by Epi- info version 7 then analyzed using SPSS for windows version 20. Descriptive analysis was done and logistic regression analyses were also used to see the association of dependent and independent variables. Finally odds ratio and 95% confidence interval were computed to determine the strength of association.

Result: This study found out practice of labour pain management methods was 43.3%, which is only non-pharmacologic methods, practice of pharmacologic method was nil. Multivariate logistic regression showed professionals positive attitude for labour pain management {AOR=2.242, 95% CI=(1.242-4.048)} and High level qualification {(AOR= 2.876, 95% CI = (1.29-6.39)} were statistically associated with practice of labour pain management methods.

Conclusion and recommendation: This study revealed poor practice of labour pain management methods. Positive attitude and highest qualification found to be significant predictor of practice of labour pain management methods. So there is a need to build attitude and knowledge of skilled attendants towards good practice of labour pain management and further studies should be done from mother satisfaction point of view.

Keywords: Labour pain; Tigray regional hospitals; Pain management; Skilled attendant

Introduction

For all woman labour involves pain, often the pain of labour is the most severe pain that a woman ever has to face. The pain a woman experiences during labour and birth is subjective, individualized and caused by a number of interrelating factors. Physical, affective, psychosocial and environmental components all shapes the pain experience [1].

Pain relief in childbirth is subject to many social and cultural modifiers, which continue to change. There is no other circumstance in which it is considered acceptable for a person to experience untreated severe pain even while under a physician's care [2]. Control of pain rather than absolute amelioration is seen by many to provide greater satisfaction. The importance of analgesia as a contribution to overall satisfaction has been recognized increasingly in the last century [3]. Sir James Simpson, professor of Obstetrics at the University of Edinburgh first advocated the use of ether for Obstetric analgesia in 1850's [4]. Queen Victoria in 1853 had the pains of childbirth relieved by intermittent whiffs of chloroform and thereafter the opposition from churches regarding obstetric analgesia remained silent [5].

Supporting women at delivery is an essential part of public health care. The term 'labor support' refers to continuous non-medical care of a laboring woman. It includes physical comforting such as touching, massaging, bathing, grooming, applying warmth or cold; and emotional support such as continuous companion, reassurance, encouragement, anticipatory guidance, information provision, and non-medical advice. Labor support and care may also involve procedures and facilitation of communication between the woman and staff to assist her in making informed choices. Furthermore, it may comprise emotional support for the woman's partner [6,7].

Labour pain management is generally classified in to two pharmacologic and non-pharmacologic labour pain management, the difference is by the use of drug for pain relief through pharmacologic pain management.

Utilization of labour pain management methods varies from place to place even it varies from one setup to another. In developed nation pharmacologic way of pain relief is the preferred one whereas in developing countries most of the time non-pharmacologic way of managing labor pain is utilized [8].

A study in Colombia reported inequity distribution of epidural analgesia in developed versus developing countries, in developed countries the use of epidural analgesia during labour is around 60% of deliveries; covering France 75%; Sweden 71% and Colombia 31.5%. In other less developed countries such as South Africa, only 21% of women used epidural [9]. In our country Ethiopia as a study conducted on all obstetric care givers in Amhara Regional State Referral Hospitals showed the overall utilization of obstetric analgesia in labour pain management was 40.1% which showed only non-pharmacologic methods. The utilization of pharmacologic obstetric analgesia methods was zero [10]. Which shows poor utilization of labour pain management methods in Ethiopia as compared to other countries, this study also report factors for poor utilization like professional's qualification and knowledge were found statistically associated with utilization of obstetric analgesia. But in general factor for utilization of labour pain management methods broadly classified in to individual factor, institutional factor and background characteristics of health professional [8].

Materials and Methods

Study design

Hospital based cross-sectional study.

Study area and period

The study was conducted at all general hospitals found in tigray region from October 01-30, 2016. Tigray region is the northernmost of the nine regions of Ethiopia. Tigray is the homeland of tigray, irob and kunama people. Tigray is also known as region 1 according to the federal constitution. Its capital is Mekelle also known as the northern star. Tigray is bordered by Eritria to the north, Sudan to the west, afar region to the east and the Amhara region to the south and southwest. According to the projected census of 2007, the region has a total population of 4,806,843 (3,787,667 rural and 1,019,176 urban). According to the 2011 report of TRHB Health care coverage in Tigray region reaches around 83%. Health care services delivered through 2 referral hospitals, 14 general hospitals, 20 primary hospitals, 214 health centers and 613 health posts. Regarding human power, currently there are 21 specialist doctors, 55 general practitioners, 2495 nurses, 1262 health extension workers and other different professionals providing preventive and curative services in the above mentioned institutions.

Sample size determination

All skilled attendants working at labour wards in all general hospitals of tigray region were taken like a census which was in number two hundred thirty three.

Sampling procedure

There are fourteen general hospitals in tigray region, which are; sebul (shire), Adigrat, Mekelle, Maychew (Ilemlem Karl), Kidist Mariam (Axum), Adwa, Meareg (dansha), Abyadi, Shiraro, Wekero, Kiha, Korem, Alamata, Humera (Kahsaye) hospital. All two hundred thirty three skilled attendants who were working at labour ward and follow laboring mother in the above general hospitals during the study period was included.

Data collection tools and procedures

Structured and pretested interviewer-administered questionnaires was prepared and utilized after reviewing relevant literatures. All skilled attendants who were working in labour wards of all the general hospitals found in tigray region and who fulfilled eligibility criteria was interviewed by structured questionnaires. Data was collected by structured interviewer-administered pre tested questionnaires and fourteen data collector was recruited one in each general hospital and four supervisor supervised the data collection.

Data quality control

Training was provided for data collectors for making common understanding of the study in general and the questioner in particular and pretest was conducted in Ayder referral hospital by delivering interviewer-administered questioner to skilled attendants to shape the questioner.

Data analysis

After data collection, the questioners were checked for completeness and code was given before data entry. Data was entered in to computer by Epi-info version 7 and exported to SPSS version 20 for further analysis. Descriptive statistics were computed to determine Frequencies and summary statistics (mean, standard deviation, and percentage) to describe the study population in relation to socio-demographic and other relevant variables. Data were presented using tables, graphs and figures. Variables with P value <0.05 in Bivariate analysis were transferred to multivariable analysis. Multivariable analysis was done to test association between predictors and dependent variables.

Ethical clearance

Ethical approval was obtained from the IRB (institutional review board) of Mekelle University, College of Health Sciences. Letter of permission was obtained from tigray regional health bureau to all general hospitals and informed consent from each study participants.

Results

Background characteristics of respondents

A total of two hundred thirty three skilled attendants were included in the study, making a response rate of 100%. The mean age of the respondents was 30.2 year, with a standard deviation of (SD ± 6.6) years, of them 152(65.3%) were in the age group of 20-29. Majority of the respondents were female 141(60.5%). Among the respondents 193 (82.8%) were orthodox Christians and 154(66.1%) were midwives. Of the respondents 81(34.8%) of were diploma holder. From the total majority of them 151(64.8%) had less than 5 years' experience (Table 1).

Table 1 Socio-demographic characteristic of skilled attendants working at labour wards in tigray region general hospitals, North Ethiopia, October 1-30, 2016 G.C (n=233).

Characteristics	Frequency	Percent (%)
Age		
20-29	152	65.3
30-39	56	24
≥ 40	25	10.7
Religion		

Orthodox	193	82.8
Protestant	21	9.1
Muslim	15	6.4
Other *	4	1.7
Profession		
Midwife	154	66.1
Medical doctor	35	15
IESO	32	13.7
Nurse	12	5.2
Highest qualification		
Diploma	81	34.8
BSc	68	29.2
MSc intern	51	21.9
Intern doctor	33	14.1
Clinical experience		
≤ 5	151	64.8
6-9	39	16.7
≥ 10	43	18.5
Other* = catholic and seventh day Adventist		

Knowledge on labour pain management method type and attitude of respondents

All respondents' 233(100%) responds as they knew about managing labour pain in general, of them 113(48.5%) knows only non-pharmacologic labour pain management methods whereas 120(51.5%) knows both pharmacologic and non-pharmacologic labour pain management methods.

Of the pharmacologic methods, systemic opioids 119(51.1%), regional analgesia 114(48.9%), non-opioid systemic analgesia 95(40.8%) and inhalational 94(40.3%) of respondents knows this methods respectively.

On the other hand; the most known non-pharmacologic labour pain management method was massage the back by all 233(100%) of respondents (Table 2).

Table 2 Professionals knowledge of non-pharmacologic method types in all TRGH, North Ethiopia, October 1-30, 2016 G.C (n=233).

Methods	frequency	percent (%)
TENS		
Yes	20	8.6
No	213	91.4
Acupuncture		
Yes	35	15
No	198	85

Massage the back		
Yes	233	100
No	0	0
Psychotherapy		
Yes	232	99.6
No	1	0.4
Show how to bear down		
Yes	229	98.3
No	4	1.7
Allow companionship		
Yes	228	97.9
No	5	2.1
Hot/cold pack immersion		
Yes	156	67
No	77	33
Breathing technique		
Yes	196	84.1
No	37	15.9
Hypnosis		
Yes	27	11.6
No	206	88.4
Divertional therapy		
Yes	100	42.9
No	133	57.1
Music therapy		
Yes	137	58.8
No	96	41.2
Subcutaneous water injection		
Yes	43	18.5
No	190	81.5

In this study out of the total respondents 140(60.1%) of professionals had adequate knowledge on list of method type while the rest 93(39.9%) respondents had inadequate knowledge even from the listed method types.

Among the respondents who knew about pharmacologic labour pain management methods 98(80%) of them reported labour analgesia had side effect by delaying progress labour and 31(25.83%) of them reported fetal distress as a side effect of labour analgesia.

Among the respondents 101(43.3%) professionals had Negative attitude while the rest 132 (56.7%) of them had Positive attitude towards managing labour pain. Among the study participants 154(66.1%) of them believed as managing labour pain help the mother on labour, but only 58(24.9%)

professionals perceive as every mother pain during labour should be managed. Among the skilled attendants 178(76.4%) of them believed that analgesia (pharmacologic method) is not necessary for managing labour pain. Of the respondents 217(93.1%) believed as labour pain is natural and the mother has to face it, but majority 177(76%) of them consider managing labour pain as their responsibility of care for laboring mother.

Personal preference and pain expectation

Among the study participants all of them, two hundred thirty three (100%) expectation of labour pain was labour pain as a severe pain. Most professionals 191(81.9%) prefer non-Pharmacologic methods to manage labour pain.

Practice of non-pharmacologic labour pain management methods

From the total of 233 skilled attendants working at all tigray region general hospitals 101(43.3%) of them practiced non-pharmacologic labour pain management methods. From the non-pharmacologic labour pain management methods, Show how to bear down was the most widely used method by 229(98.3%) professionals followed by psychotherapy 179(76.8%) and allow companionship 135(57.9%) (Figure 1).




Figure 1 Proportion of utilization of non-pharmacologic labour pain management methods by skilled attendants working in general hospitals of tigray region, North Ethiopia, October 1-30, 2016 G.C.

Reason for non-utilization

The most common reasons mentioned by respondents for non-utilization of labour pain management were high patient flow by 233(100%) of respondents and small number of staff by 177(76%) of respondents (Figure 2).




Figure 2 Reason given by skilled attendants for non-utilization of labour pain management in TRGH, North Ethiopia October 1-30, 2016 G.C.

Factor associated with practice of non-pharmacologic labour pain management methods: On the bivariate analysis professional age, level of knowledge, attitude, profession and professional's highest qualification was identified to be significantly associated with current practice of labour pain management methods. But only attitude and highest qualification were remained significantly associated with current practice of non-pharmacologic labour pain management methods in the multiple logistic analysis.

The odds of current practice of non-pharmacologic labour pain management methods was higher among professionals who have positive attitude for managing labour pain. Skilled attendants who have positive attitude for managing labour pain were about 2.24 times more likely to practice non-pharmacologic labour pain management methods as compared to those professionals who have negative attitude for labour pain management {AOR=2.242, 95% CI= (1.242-4.048)}.

Professionals who had highest qualification of MSc (high level qualification) were 2.876 times more likely to practice non-pharmacologic labour pain management methods than professionals who had lower level qualification of diploma. {(AOR=2.876, 95% CI = (1.29-6.39))} (Table 3).

Institutional factor

From the total respondents one hundred eighty one (77.7%) reported as allowing companion of choice it's not allowed by their hospital and still all respondents didn't gate any special training on managing labour pain. From the respondents who knows pharmacologic method type ninety six (80%) of them reported as Pethidine is available in their hospital.

Table 3 Bivariate and multivariate analysis of factors associated with practice of non-pharmacologic labour pain management methods in TRGH, North Ethiopia, 2016.

	Practice of non-pharmacologic labour pain management method		COR (95% CI)	AOR (95%CI)
	Yes	No		
	Frequency (n)	Frequency (n)		
Age				
20-29	70	82	2.70(1.02-7.14)	2.25(0.74-6.83)
30-39	25	31	2.55(0.88-7.36)	1.37(0.44-4.26)
≥ 40	6	19	1	1
Qualification				
Lower level	26	55	1	1
Medium level	41	60	1.45(0.78-2.66)	1.28(0.68-2.40)
Higher level	34	17	4.23(2.01-8.92)	2.87(1.29-6.39)**
Attitude				
Positive Negative	72	60	2.98(1.72-5.17)	2.24(1.24-4.05)**
	29	72	1	1
Knowledge				
Adequate	73	67	2.53(1.45-4.34)	1.34(0.67-2.67)
Inadequate	28	65	1	1

Profession				
Midwife	59	95	0.55(0.31-0.95)	0.96(0.51-1.81)
Other	42	37	1	1
**= significant variable in backward stepwise logistic regression				
Lower level: diploma, Mid-level: BSc and medical intern, Higher level: MSc				
Other: Nurse, Medical intern, IESO				

Discussion

There is no doubt that provision of pain relief services in labour would add value to obstetric services, managing labour pain is not luxury rather anticipated obstetrics care [11]. This institutional based cross sectional study has attempted to determine utilization of labour pain management methods and associated factors by skilled attendants working in tigray region general hospitals at labour ward, North Ethiopia.

Childbirth is one of the most painful events that a woman is likely to experience, the multidimensional aspects and intensity of labour pain is not comparable with other circumstances, but majority of the respondents in this study 178(76.4%) of them believed that analgesia (pharmacologic method) is not necessary for managing labour pain, which is unlike the study in Zaria, Nigeria. There 94.8% of health care provider agreed that labour analgesia is needed during labor [11]. The possible reason for this difference might be the awareness of pharmacologic agents used to relieve pain during labor by health care provider in Zaria, Nigeria was high as compared with this study, which is observed by the attitude difference between these two populations.

The current study found out that the proportion of skilled attendants utilizing non-pharmacologic labour pain management methods were 43.3%. This study finding is in line with the study done in Bangladesh; there utilization of non-pharmacologic labour pain management was 40%. But specific to each method utilization, in this study utilization of non-pharmacologic labour pain management methods was lower than Bangladesh. While comparing utilization in Tigray, Ethiopia versus Bangladesh, labour exercise 56.2% versus 78.4%, massage the back 47.6% versus 53.6%, breathing technique 28.3% versus 40.2% and allow companionship 57.9% versus 77.3% of respondents' utilization. But higher than Bangladesh by showing the mother how to bear down 98.3% versus 72.2% of respondents utilization [5]. This difference is expected difference from place to place because health professionals couldn't be identical in every aspect from one site to another.

Regarding pharmacologic labour pain management this study found out proportion of skilled attendants utilizing pharmacological labour pain management methods was zero, the finding of this study is in line with a study done in Amhara region referral hospital which was zero [10]. But this study finding is much lower as compared to other studies like obstetric analgesia utilization in Addis Ababa, Ethiopia (54.2%) [12], epidural obstetric analgesia use in France (75%), Sweden

(71%), Bangladesh (58.5%) [5], Colombia (31.5%) and South Africa (49%) [9]. Even in Nigeria 49% of obstetricians offered obstetric analgesia of which the commonest analgesia was opioids (41.1%)(13). And 27% of mothers in Canada received the commonest (92.6%) IM Pentazocine Hydrochloride for labour pain relief [13,14]. The possible reason for this incomparable difference is this study was conducted in tigray region general hospitals, they are found in rural area they are far from the capital Addis Ababa, but in other countries the hospitals are modern and all equipment's and drugs for labour analgesia are available in those modern setup. And the other possible reason is vast majority of the respondent's here are diploma holder and even they didn't have adequate knowledge about pharmacologic labour analgesia and the other reason might be negative attitude towards utilization of labour analgesia.

This study also revealed those skilled attendants who have positive attitude for managing labour pain were 2.24 times more likely to use labour pain management method than skilled attendants who have negative attitude. {AOR(95%CI)=2.24(1.24-4.05)} which is consistent with a study in Bangladesh, which reported negative attitude of health care providers were restrictive towards utilization of analgesic for normal labour, and they also thought it would interfere with normal progress of labour. However, they expressed positive attitude to help mothers to cope with labour pain through assurance and carry out some activities to comfort the women in labour [5].

In this study highest qualification was statistically significant predictor of utilization of labour pain management methods. Skilled attendants who were MSc intern (higher level qualification) were 2.87 times more likely to use labour pain management methods than professionals who had diploma (low level qualification). {AOR (95%CI)=2.87(1.29-6.39)}.

This finding is opposite with study done in Amhara regional state referral hospitals which reported Obstetric care givers who had diploma (low level qualification) were 2.69 times more likely to use obstetric analgesia methods than professionals who had second degree (high level qualification) {AOR(95%CI)=2.69(1.13-6.41)} [10].

The possible explanation for this difference might be in this study professionals who have higher level qualification are MSc interns, since they are students so their evaluation also depend on their management skill and client professionals' relationship, so this implication is reflected by their utilization of labour pain management methods.

In this study professionals reason for not using labour pain management method were; high patient flow, small number of staff, lack of knowledge and skill and unavailability of equipment's and drugs for managing labour pain. The finding is consistent with a study done at public hospitals of Addis Ababa Ethiopia [12].

Limitation and Strength of the Study

Limitation of the study

In this study head nurses were recruited as a data collector which has social desirability bias.

Strength of the study

In this study census was employed to recruit study participants, which eliminate sampling error.

Conclusion

Even though managing labour pain is anticipated obstetrics care, this study revealed poor practice of non-pharmacologic labour pain management methods. The major predictor for practice of non-pharmacologic labour pain management methods as per this study finding was professionals' positive attitude towards managing labour pain and highest qualification.

In a setup like us with high patient flow practicing methods for managing labour pain was considered as luxury rather than anticipated obstetrics care, as a result high patient flow was the number one listed reason for non-practicing of labour pain management methods.

Recommendation

Federal ministry of health and TRHB

To incorporate education of pharmacologic and non-pharmacologic labour pain management methods in the curriculum at all level of qualification including diploma program.

Tigray region general hospitals

An institutional framework that includes education and supply of required equipment will help care providers to bridge the gap between their attitudes and provision of pain relief services in labor.

Non-governmental organization

To participate in education, positive attitude creation and information regarding benefits of managing labour pain and strengthen short term trainings on issues related to labour pain and obstetric analgesia.

Researcher

A qualitative study needs to be done on utilization of labour pain management methods from mothers' satisfaction point of view.

References

1. Beigi NM, Broumandfar K, Bahadoran P, Abedi HA (2010) Women's experience of pain during childbirth. *Iran J Nurs Midwifery Res* 15: 77-82.
2. Eisenach JC, Pan P, Smiley RM, Lavand'homme P, Landau R, et al. (2013) Resolution of pain after childbirth. *Anesthesiology* 118: 143-151.
3. Soet JE, Brack GA, Dilorio C (2013) Prevalence and predictors of women's experience of psychological trauma during childbirth. *Birth (Berkeley, Calif)* 30: 36-46.
4. Lawrence C (1999) Medicine's 10 greatest discoveries. *BMJ* 319: 389.
5. Tasnim DS (2010) Perception about pain relief during normal labour among health care providers conducting delivery. *Medicine today* 22: 20-23.
6. Bonnet MP, Prunet C, Baillard C, Kpea L, Blondel B, et al. (2017) Anesthetic and obstetrical factors associated with the effectiveness of epidural analgesia for labor pain relief: An observational population-based study. *Reg Anesth Pain Med* 42: 109-116.
7. Klein MC, Grzybowski S, Harris S, Liston R, Spence A, et al. (2001) Epidural analgesia use as a marker for physician approach to birth: Implications for maternal and newborn outcomes. *Birth (Berkeley, Calif)* 28: 243-248.
8. Mugambe JMNM, Hiemstra LA, Steinberg WJ (2007) Knowledge of and attitude towards pain relief during labour of women attending the antenatal clinic of Cecilia Makiwane Hospital, South Africa. *Medpharm* 49: 16-20.
9. Ortiz GD, Navarro-Vargas JR, Eslava-Schmalbach J (2013) Inequity in healthcare-the outlook for obstetric analgesia. *Colombian J Anesthesiology* 41: 5-7.
10. Bitew AWA, Seyum T (2016) Utilization of obstetric analgesia in labor pain management and associated factors. *J Biomedical Sci* 5: 1-6.
11. Ogboli-Nwasor E, Adaji S, Bature S, Shittu O (2011) Pain relief in labor: A survey of awareness, attitude, and practice of health care providers in Zaria, Nigeria. *J Pain Res* 4: 227-232.
12. Mulugeta H (2016) The practice of labor analgesia and its perceived barriers among health care providers working in public hospitals of Addis Ababa, Ethiopia.
13. Lawani JNELO, Okechukwu BA, Iyoke CA, Ekem NN (2014) Obstetric analgesia for vaginal birth in contemporary obstetrics: A survey of the practice of obstetricians in Nigeria. *BMC* 14: 1-6.
14. Likis FE, Andrews JC, Collins MR, Lewis RM, Seroogy JJ, et al. (2014) Nitrous oxide for the management of labor pain: A systematic review. *Anesth Analg* 118: 153-167.